



Stability Constants of Ternary Complexes of Transition Metal(II) ions with Aspartic Acid and Glutamic Acid as Primary Ligands and Ampicillin and Cephalexin as Secondary Ligands

A.B. PATIL

Department of Chemistry, P.N. College, Pusad - 445 216, India.

*Corresponding author E-mail: arunpatil2012@rediff.com

(Received: March 20, 2012; Accepted: April 18, 2012)

ABSTRACT

The interaction of Mn (II), Co (II), Ni (II), Cu (II) and Zn (II) metal ions with ampicillin (AMP) and cephalexin (CEP) have been studied by pH-metric technique at 0.1 M (KNO₃) ionic strength at 302 ± 0.5 K in aqueous medium. The data obtained were used to evaluate the values of proton-ligand and metal-ligand stability constants using Irving-Rossotti titration technique. Mixed ligand complex studies of these metal ions using aspartic acid (ASP) and glutamic acid (GLU) as primary ligands and AMP and CEP as secondary ligands also have been carried out pH-metrically at the same conditions.

Key words: Ternary complexes, Transition metal ions, ampicillin, aspartic acid and glutamic acid.

INTRODUCTION

Recently, there has been considerable interest in the study of binary, ternary and quaternary complexes by pH-metric method¹⁻³. The ligand ampicillin (AMP) and cephalexin (CEP) are well known antibiotics⁴⁻⁶.

Ternary complexes of Ni (II) and Cu (II) with nicotinic acid as primary ligand and imidazoles, benzimidazole, histamine and L-histidine as secondary ligands have been studied by Nair and Neekantan⁷. Patil and Mhaske⁸ and Job's Turner and Andersom and Iqbal *et al.*,¹⁰ have studied the stability constants of Mn (II), Co (II), Ni (II), Cu (II)

and Zn (II) with nitrilotriacetic acid and Iminodiacetic acid as primary ligands and ampicillin and cephalexin as secondary ligands potentiometrically. Nigam and coworkers¹¹ have studied the ternary complexes of Mn (II), Co (II), Ni (II), Cu (II) and Zn (II) using ASP as primary ligand and thymine as secondary ligand potentiometrically. Ternary complexes of Cu (II) using ASP and GLU as primary ligands have been reported potentiometrically by Pandeya and Patel¹¹. Kalshetti and coworkers¹¹ have studied binary and ternary complexes of Co (II), Ni (II), Cu (II) and Zn (II). Mixed ligand complexes of transition metal(II) ions with N-(2-hydroxybenzylidene)-2,3-dimethylaniline as primary ligand and N-(2-hydroxy-1-naphthylidene)-

4-chloroaniline as secondary ligand has been studied by Mapari and Mangaonkar¹³. Solution equilibria of ternary systems involving transition metal ions, hydroxamic acids, and bioligands have been studied by Khalil and Mahmoud¹⁵.

In this paper the stability constants of ternary complexes of Mn (II), Co (II), Ni (II), Cu (II) and Zn (II) ions with ASP and GLU as primary ligands and AMP and CEP as secondary ligands at 302 ± 0.5 K and at fixed ionic strength, $\mu = 0.1\text{M}$ KNO_3 using modified form of Irving-Rossotti pH-metric technique¹⁶ in aqueous medium have been studied.

EXPERIMENTAL

The ligand AMP was obtained from Zim laboratories, Kalmeshwar (Nagpur) and CEP was of Relief Laboratories Kalmeshwar (Nagpur). These ligands were used as such. Carbonate free sodium hydroxide solution was prepared by standard method¹⁵. All other solutions were prepared in doubly distilled water.

The pH-metric measurements were carried out by using Elico digital pH-meter model L-120 with combined glass-calomel electrode with an accuracy of ± 0.01 of pH unit at 302 ± 0.5 K. The pH-meter was standardized against 0.05 M potassium hydrogen phthalate solution in acid medium and 0.01M borax solution in alkaline medium.

For determination of proton-ligand stability constant of the secondary ligand and the metal-ligand stability constants of ternary complexes, the following set of solutions were prepared and titrated against standard alkali solution.

Ternary Systems

- i) $9.6 \times 10^{-3}\text{M}$ HNO_3
- ii) $9.6 \times 10^{-3}\text{M}$ HNO_3 + $1.0 \times 10^{-3}\text{M}$ secondary ligand
- iii) $9.6 \times 10^{-3}\text{M}$ HNO_3 + $1.0 \times 10^{-3}\text{M}$ primary ligand + $1.0 \times 10^{-3}\text{M}$ metal ion
- iv) $9.6 \times 10^{-3}\text{M}$ HNO_3 + $1.0 \times 10^{-3}\text{M}$ primary ligand + $1.0 \times 10^{-3}\text{M}$ metal ion + $1.0 \times 10^{-3}\text{M}$ secondary ligand

The ionic strength was maintained constant (0.1M) by adding required volume of 1M KNO_3 . The ratio of metal (M): primary ligand (A): secondary ligand (L) was maintained at 1: 1: 1 in each of the ternary system.

RESULTS AND DISCUSSION

Proton-Ligand Stability Constants

The plots of volume of alkali (NaOH) against pH-meter readings were used to evaluate the proton-ligand stability constants of AMP and CEP. The deviation between free acid titration curve and secondary ligand titration curve was used to evaluate the formation functions \bar{n}_A . The proton-ligand formation curves were then obtained by plotting the values of \bar{n}_A versus pH-meter readings. From the graphs, the values of $\log K^H$ were evaluated by half integral method (method A) and point wise calculation method (method B) and presented in Table 1.

Metal – Ligand Stability Constants of Ternary Complexes

^{n_A} The metal – ligand stability constants of the ternary complexes were evaluated assuming that the formation of hydrolyzed products, polynuclear complexes, hydrogen and hydrogen bearing complexes were absent. An examination of titration curves indicates that complex formation has taken place in the solution on the following grounds:

1. The ternary complex titration curves show the displacement with primary complex titration curves. The horizontal distance was measured between acid curve and the secondary ligand curve ($V_2 - V_1$) and subtracted through the horizontal distance between ternary complex curves and primary

Table 1: Proton-Ligand Stability Constants of AMP and CEP

Ligand	$\log K^H$	
	Method A	Method B
AMP	7.35	7.37
CEP	7.40	7.42

Table 2: Metal-Ligand Stability Constants of Ternary Complexes


System	$\log K_{ML}^{MA}$									
	Mn (II)		Co (II)		Ni (II)		Cu (II)		Zn (II)	
	A	B	A	B	A	B	A	B	A	B
[M (II) - (ASP) (AMP)]	3.79	3.78	3.97	3.96	4.14	4.15	4.62	4.64	3.90	3.91
[M (II) - (GLU) (AMP)]	3.60	3.61	3.67	3.69	4.06	4.05	4.59	4.59	3.81	3.82
[M (II) - (ASP) (CEP)]	3.87	3.88	4.12	4.13	4.42	4.43	5.28	4.29	4.30	4.29
[M (II) - (GLU) (CEP)]	3.74	3.75	3.94	3.94	4.19	4.18	5.24	5.23	4.06	4.05

complex titration curves ($V_4 - V_3$) show the positive difference which proves the earlier released of protons in the formation of ternary complexes.

- The hydrolysis of the metal ions was suppressed and precipitation did not result.

The values of \bar{n} vary from 0 to 0.75, thus confirming the formation of 1:1:1 mixed ligand complexes. The values of $\log K_{ML}^{AS}$ and $\log K_{ML}^{GLU}$ have been evaluated from the formation curves (vs. pL). At $\eta = 0.5$ in the formation curve, $pL = \log K$. The metal- ligand stability constant of AMP as secondary ligand and ASP and GLU as primary ligand are presented in Table 2.

The Irving-Williams order^{16, 17} of stability constants was observed in ternary complexes which is



This sequence of stability of complexes with respect to metal ion is due to decreasing atomic radius and increasing the second ionization potential.

ACKNOWLEDGMENTS

The authors are thankful to UGC and Principal, P. N, College, Pusad for providing necessary facilities.

REFERENCES

- A. E. Martell and R. M. Smith, *Critical Constants-1: Amino acids*, New York (1974).
- H. Sigel, *Metal Ions in Biological Systems-2*, Marcel-Dekker, Inc., New York (1973).
- M. T. Beck, *Chemistry of Complex Equilibria*, Van Nostrand, New York, pp.174-190 (1970).
- The Pharmaceutical Codex, 11th Edition, The Pharmaceutical Press, London (1979).
- Clarke's Isolation and Identification of Drugs, 2nd Edition, The Pharmaceutical Press, London (1986).
- Remington's Pharmaceutical Science, 17th Edition, Mack Publishing Co., Easton, Pennsylvania (1985).
- M.S. Nair and M. A. Neelkantan, *Indian J. Chem.*, **38A**, 1307(1999).
- A. B. Patil and T. H. Mhaske, *Orient J. Chem.*, **17**(3): 483-486 (2001).
- S.A. Iqbal, Sibi Jose and Ishaq Zaafarany, *Orient. J. Chem.* **28**(1): 613-618 (2012).
- Turner S.E. and Anderson R.C., *Chem. Soc. J. Am.* **7**(1): 912 (1949).
- N. B. Nigam, P. C. Sinha, M. Gupta and M. N. Srivastava, *Indian J. Chem.*, **22A**: 818 (1983).
- K.B. Pandeya and R. N. Patel, *Indian J. Chem.*, **30A**, 193 (1991).
- B.M.Kalshetty, S.S.Suvarnakhandi, B.I.Karlatti and M.B.Kalashetti, *International Journal of Applied Biology and*

- Pharmaceutical Technology*, **2**(2): 386 (2011).
14. A. K. Mapari and K. V. Mangaonkar, *International Journal of Chem Tech Research*, **3**(2): 802-805 (2011).
 15. M. M. Khalil and R. K. Mahmoud, *J. Chem. Eng. Data*, **52**(2): pp789-797(2010).
 16. H.Irving and H. S. Rossotti, *J. Chem. Soc.*, **76**, 2904(1954).
 17. A.I. Vogel, *A Text Book of Quantitative Analysis*, Longman, London, pp.241 (1961).
 18. H.Irving and R.P.J. Williams, *Nature*, **162**: 746 (1948)
 19. H.Irving and R.P.J. Williams, *J. Chem. Soc.*, **76**: 3192 (1953).