



## Flavonoids Of *Astragalus alopecua* Ðàll

**A.K.PATSAYEV\*, B.K.MAKHATOV,  
A.Y.BUKHARBAYEVA and K.DZH.KUCHERBAYEV**

South-Kazakhstan state pharmaceutical academy, Republic of Kazakhstan, Shymkent

\*Corresponding author E-mail: anapia.patsaev@mail.ru

<http://dx.doi.org/10.13005/ojc/330353>

(Received: March 16, 2017; Accepted: May 24, 2017)

### ABSTRACT

*Astragalus alopecua* is one of the valuable traditional medicinal herbs that exhibits a line of biological activities. In this research, the content of *Astragalus alopecuas* flavonoids were studied. From the generative and vegetative organs *astragalus* identified a number of active compounds (flavonoids, polysaccharides, triterpene saponins). We conducted preliminary study the aerial part of *Astragalus alopecuas* showed the presence of the most common groups of biologically active substances, flavonoids, triterpenoids. We identified quantitative content of flavonoids by UV-spectrophotometry in conversion to quercetin from aerial parts, in flowers and in root of plant *Astragalus alopecua*. As a result we found that in leaves and stems appeared highly accumulation of flavonoids. Contents of total flavonoid reached up to 18% and the major ingredient is quercetin.

**Keywords:** *astragalus alopecua*, spectrophotometry, flavonoids

### INTRODUCTION

One of the perspective sources of phyto preparations are medicinal plants containing flavonoids, which is by the wide spread in plants and higher structural diversity now a days is in the center of researchers attention in the field of pharmacy and medicine (Korulkin et al., 2007; Kurkin, 2002; Kurkin, 2007; Makarova and Makarov, 2010; Muraviyeva, 2002; Flavonoids: Chemistry, Biochemistry, and Applications, 2006; The Flavonoids: Advances in Research, 1982; Wagner, 1993). Flavonoids – are more class of natural phenol compounds, which characterized by structural multiplicity, higher

various activity. Widely amplitude of biological activity of flavonoids are connected with diversity of their chemical structures and different physical-chemical properties. This interest is connected with circumstance, that flavonoids, which is being evolutionary adequate to human organism, stipulates anti-oxidative, vasoprotective, hepatoprotective, cholagogue, diuretic, neurotropic actions (Korulkin et al., 2007; Kurkin, 2002, 2007; Kurkin and Pravdiceva, 2008; Kurkin, 2009). One of the species *Asrtagalus membranaceus* has been used for more than 2,000 years in China to boost the body's general vitality and strengthening resistance to exogenous pathogens (Fangli Lu, 2014).

Flavonoids variety distributed in nature. As the food-derived cancer preventive compounds, they display potential therapeutic benefit in cancer and may be considered as the candidates for chemotherapeutic agents (Sak, 2014). To improve their biological activities and enhance their anticancer effects, it is also important to chemically modify the structures of natural flavonoids by introducing structural variations into their backbone (Xiaofeng Shi *et al.*, 2016).

The most reported activity of flavonoids is their protection against oxidative/nitrosative stress. Thus flavonoids can scavenge peroxy radicals, and are effective inhibitors of lipid peroxidation, and can chelate redox-active metals, and therefore prevent catalytic breakdown of hydrogen peroxide (Ergul Belge Kurutas, 2016). Generally, the synthesis of flavonoids is enhanced when the plants face with stress agents, such as strong light and there have been few reports on the MS fragmentation of them (Wang Hui, 2015). Around 840 new isoflavonoids have been reported over last 15 years indicating the increasing interest in search for the natural isoflavonoids (Sharma, 2013).

In terms of the species number, *Astragalus* L. (Fabaceae), represented by about 10 subgenera, 130 sections, and nearly 2500 taxa, is the richest genus of vascular plants on earth. Erect, caulescent, usually stout, perennial herbs. Stipules free from the petiole, rarely decurrent, linear, lanceolate, linear to lanceolate, triangular, triangular lanceolate, ovate to acuminate, glabrous or pilose, ciliate. Leaves imparipinnate (Hasan Akan and Zeki Ayta, 2014).

More than 40 years ago, research on *Astragalus* species was initiated in Bulgaria focused on numerous biologically active metabolites. Novel and unusual for the genus, group of compounds named flavoalkaloids were identified in *A. monspessulanus* subsp. *monspessulanus*. Flavoalkaloids were previously known only as aglycones (Krasteva *et al.*, 2016).

Four new flavonoids, astragaisoflavans, along with thirteen known ones were isolated from the aerial parts of *Astragalus hoantchy*. Compound 1 and 4 were evaluated for their antifungal activity against *Alternaria solani* (Kai Guo *et al.*, 2016).

As a consequence we carried out *Astragalus alopecica* Pall assessment of flavonoids compounds content. The goal of the work is conclude of quantitative determination of flavonoids in plant *Astragalus alopecica* Pall.

#### Materials and methods

Object of investigation is air-dry aerial part, flower and leaves of plant *Astragalus alopecias* Pall., harvested in 2015 in South-Kazakhstan oblast (Ordabassy region, Kozhatogai village) during blossoming period of the plant and roots collected in fruiting period. To avoid any destruction of chemical components, the collected materials were dried in the shade. The plant was identified by candidate of biological sciences T.S.Ibragimov from South-West scientific-industrial Center of agricultural holdings.

UV spectra were recorded using spectrophotometer SPh-2000 by LOMO. For determination of flavonoids is used in complex formation reaction with aluminum chloride solution exclude the deposit in account optical density of other group of compounds in conversion to quercitin (Muraviyeva *et al.*, 2002).

Analytical sample of raw material crushed to particles, pervaded through sieve with diameter ports 1 mm. About 1g (precisely weighted amount) raw material put in the flask with volume 150 ml added 30 ml 90% ethanol, containing 1% concentrated hydrochloric acid, flask connected to diverse refrigerator and heated on boiling water bath during 30 minutes. Then flask cooled till room temperature and filtered through paper filter in volumetric flask with volume 100 ml. Extraction is repeated twice. Then extracts put together, filtered through same filter to the same flask, washed filter with 90% ethanol and add it till mark (A solution).

In volumetric flask with volume 25 ml put 2 ml A solution, add 1 ml 1% aluminum chloride solution in 95% ethanol add 95% ethanol till mark. After 20 minutes measured the optical density of solution on spectrophotometry at 430 nm wave length in 10 mm layer thickness cuvet.

As a blank solution used solution containing 2 ml A solution, add 95% ethanol till mark in 25 ml volumetric flask.

**Table 1**

<b>Astragalus alopecias</b>	<b>Leaves</b>	<b>Flower</b>	<b>Root</b>
	18%	2,8%	2,0%

**RESULTS**

Content of flavonoids in conversion to quercetine and absolute dry raw material in percentage ( $\bar{O}$ ) is calculated by the formula:

$$X = \frac{D \cdot 25 \cdot 100 \cdot 100 \cdot 100}{764.6 \cdot m \cdot 2 \cdot (100 - W)}$$

where,

D – optical density of investigated solution at wave length 430 nm;

764.6 – indicator of absorbance of quercetin complex with 1% aluminum chloride solution at wave length 430 nm;

m – mass of raw materials in g;

W – mass lost while drying.

Content of flavones in conversion with quercetin

Obtained results may suggest the presence of anti-oxydative activity of plant *Astragalus alopecia* Pall.

**CONCLUSIONS**

Flavonoids have the main significance in treatment of different disease. One of the flavonoid containing plant is a *Astragalus alopecia* show interest for studying. Study of chemical composition of astragals is show presence of flavonoids, phenylcarbonic acids, oxycorich acids, polysaccharides and other compounds. Pharmacological investigations show, that one of the group of acting substances equal with polysaccharides are phenol compounds, and in particular, flavones.

By the UV- spectrophotometry method was determined the quantitative content of phenolic compounds in plant *Astragalus alopecia* Pall. Results of investigations can be used on making determined tarts of analytical documents on medicinal plant raw materials and to form new phytoproducts.

**REFERENCES**

1. Ergul Belge Kurutas, The importance of antioxidants which play the role in cellular response against oxidative/nitrosative stress: current state, *Kurutas Nutrition Journal*, , **2016**, *15*,. 71.
2. Fangli Lu, Treatment Methods of Traditional Chinese Medicine for Toxoplasmosis, Treatment of Human Parasitosis in Traditional Chinese Medicine, Parasitology Research Monographs *Springer-Verlag Berlin Heidelberg*, **2014**. 6.
3. Flavonoids: Chemistry, Biochemistry, and Applications, Ed. by Andersen ØM and Markham KR, Boca Raton; London; New York: CRC Press Taylor & Francis Group, **2006**, 1197.
4. Hasan Akan, Zeki Ayta, The revision of the section Alopecuroidei of the genus *Astragalus* (Fabaceae) in Turkey, *Turkish Journal of Botany, TUBÝTAK*, , **2014**, *38*,37-59.
5. Kai Guo, Xiaofeng He, Yanping Zhang, Xiuzhuang Li, Zhiqiang Yan, Le Pana, Bo Qina, Flavonoids from aerial parts of *Astragalus hoantchy*, *Fitoterapia*, **2016**, *114*, , 34–39.
6. Korulkin DY, Abilov ZhA, Muzychkina RA, Tolstikov GA, Natural Flavonoids, Novosibirsk: Scholar publisher «Geo», **2007**, 232.
7. Krasteva I, Shkondrov A, Ionkova I, Zdraveva P, Advances in phytochemistry, pharmacology and biotechnology of Bulgarian *Astragalus* species, *Phytochem Review, Springer Science, Business Media Dordrecht*, **2016**.
8. Kurkin VA, Flavonoids of Pharmacopeia plants, Samara: SamSMU; LTD «Ophort», **2012**, 290.
9. Kurkin VA, Fundamentals of phytoterapy: Textbook for pharmaceutical faculty students, Samara: SamSMU; LTD «Ophort», **2009**, 963

10. Kurkin VA, Modern aspects of chemical classifications of biological active compounds of medicinal plants, *Pharmacy*, **2002** 50(2), , 8–16.
11. Kurkin VA, Pharmacognosy: study book for pharmaceutical faculty students, Edn 2, Samara: LTD «Ophort»; «SamSMU Russian health care», **2007**, 1239.
12. Kurkin VA, Pravdiceva OY, Saint John's flower: results and perspectives of creation medicinal substances, Samara: SamSMU; LTD «Ophort», **2008**, 127
13. Makarova MN, Makarov VG, Molecular biology of flavonoids: reference book for doctors, **2010**, 428.
14. Muraviyeva DA, Samylina IA, Yakovlev GP, Pharmacognosy: textbook, Ì.: Medicine, **2002**, 656.
15. Sak K, Cytotoxicity of dietary flavonoids on different human cancer types, *Pharmacogn Rev.*, 8(16), 2014, 122–46.
16. Sharma V, Ramawat KG, Isoflavonoids, Natural Products», Springer, Berlin, 2013, 1849–1865.
17. The Flavonoids: Advances in Research. Ed. by Harborne JB, Mabry TJ, London; New York: Chapman and Hall., **1982**, 744
18. Wagner H, Pharmazeutische Biologie. Drogen und ihre Inhaltsstoffe, Stuttgart-New York: Gustav Fischer Verlag, **1993**, 522.
19. Wang Hui, Wang Jiayuan, Wu Xiaopeng, Shen Haiyan, Luo Ying, Dai Haofu and Mei Wenli, HPLC-ESI-MS Analysis of Flavonoids Obtained from Tissue Culture of *Dracaena cambodiana*, *Chem. Res. Chin. Univ.*, **2015.**,31(1),38-43.
20. Xiaofeng Shi, Dongyan Liu, Junmin Zhang, Pengbin Hu, Wei Shen, Bin Fan, Quhuan Ma and Xindi Wang, Extraction and purification of total flavonoids from pine needles of *Cedrus deodara* contribute to anti-tumor in vitro, *BMC Complementary and Alternative Medicine*, **2016**.