



Investigation of Anticancer Agents from the Bark of *Gyrinops versteegii* (GILG.) Domke from Lombok Island

SURYA HADI^{1*}, DIAN LESTARI¹, NI KOMANG TRI DHARMAYANI¹,
BAIQ DESY RATNASARI², M ITO³, IYAMADA⁴ and TRI MULYANINGSIH⁵

¹Department of Chemistry, The University of Mataram, Mataram-83126, Indonesia.

²Department of Pharmacy, Stikes Kusuma Bangsa Mataram-83126, Indonesia.

³Graduate School of Pharmaceutical Sciences, Kyoto University, Japan.

⁴Center for Southeast Asian Studies, Kyoto University, Kyoto-606-8501, Japan

⁵Department of Biology, The University of Mataram, Mataram-83126, Indonesia.

*Corresponding author E-mail: sur_hadi88@yahoo.com

<http://dx.doi.org/10.13005/ojc/360605>

(Received: October 20, 2020; Accepted: November 25, 2020)

ABSTRACT

This study is part of a research collaboration between Kyoto University and The University of Mataram, aiming to identify and to utilize the agarwood plants, i.e., *Gyrinops versteegii* from Indonesia. The study, specifically, aims to discover anticancer agents from the bark of *G. versteegii* from Lombok Island of Indonesia. There were three provenances of *G. versteegii* observed, namely Soyun, Pantai, and Buaya. Based on the Brine Shrimp Lethality Test (BSLT), *G. versteegii* Buaya showed the highest cytotoxicity with LC₅₀ 45,94 µg/ml. Meanwhile, *G. versteegii* Soyun and Pantai have LC₅₀ 75.86 µg/mL and 56.36 µg/mL respectively. A phytochemical study showed that the methanol extract of *G. versteegii* Buaya containing compounds from the group of alkaloid, flavonoid, steroid, triterpenoid, and saponin. The methanol extracts fractionated using Vacuum Liquid Chromatography into 10 fractions (F1-F10) and then retested BSLT. Among the fractions, F2 showed the best potency as an anticancer with LC₅₀ 64,12 µg/mL. Based on the GC-MS analysis, the cytotoxicity from both methanol extract and F2 is predicted to be influenced by the same compounds, namely, 1,4-Benzenediol, 2-methyl, Pyridoxylamine, 2,3-Dimethylhydroquinone, Tetramethyl-p-benzoquinone and Benzofuran. Overall the bark of *G. versteegii* from Lombok Island has great potency as an anticancer.

Keywords: Agarwood, Anticancer, *Gyrinops versteegii*, Lombok Island, Provenance.

INTRODUCTION

In an attempt to preserve the germplasm of agarwood, especially in the West Nusa Tenggara, a collection and an identification of agarwood plant is continuously conducted. *Gyrinops versteegii* (Gilg.) Domke is one of the agarwood species found in Nusa

Tenggara, specifically Lombok Island. This species is classified as superior agarwood that produces high-quality resin. As a result, the demand for *G. versteegii* is increasing by the year. Thus, exploration and plantation of this species are vigorously conducted, specifically in Lombok's forest.¹⁻³ *G. versteegii* in West Lombok is found in five provenances, namely



"*G. versteegii Buaya*"; "*G. versteegii Soyun*"; "*G. versteegii Pantai*"; "*G. versteegii Madu*"; and "*G. versteegii Beringin*".^{2,4}

The aromatic resin of the agarwood is the most attractive part of the plant. It is processed as fragrant oil to produce perfume and medicine. The other parts of the plant such as the leaves, the barks, the fruits, and the flowers are not fully recovered industrially and scientifically. Nevertheless, they possess many valuable bioactivities to maintain health and to treat some ailments like cirrhosis, tumor and cancer.⁴ However a clinical study of those bioactivities has not been conducted widely, especially in Lombok Island. Therefore the total utilization of agarwood parts, especially the barks, not only the resin part, is eagerly needed.

Hadi *et al.*,⁵ pronounced the major compound of *G. versteegii* bark a flavonoid. Flavonoid is well known to possess an anti-cancer activity. However the study has not been conducted widely. Generally, agarwood plants have good bioactivity as anti-cancer. Another agarwood species like *Aquilaria malaccensis* was reported to inhibit the proliferation of HCT116 with IC₅₀ 4 µg/mL.⁶ Besides, it also effective against lymphocytic leukemia cell P-388 with ED50 0,35 µg/mL.⁷ Further study found that the secondary metabolites, 2-(2-phenylethyl) chromones, that was isolated from agarwood was showing anti-cancer activity with IC₅₀ 14,6 µg/mL against SGC-7901.⁸ Thus, theoretically, the species of *Gyrinops* has the same potential as the *Aquilaria*'s as an anti-cancer agent.

This research is designed to investigate the compound that is responsible for the anti-cancer in the bark of *G. versteegii* of Lombok Island from three provenances (*G. versteegii Soyun*, *G. versteegii Pantai* and *G. versteegii Buaya*) using Brine Shrimp Lethality Test (BSLT). Furthermore, the chemical compound in the barks was identified based on Gas Chromatography-Mass Spectrometry (GC-MS).

MATERIAL AND METHODS

Preparation

The bark samples were obtained from three provenances of *G. versteegii* (*Soyun*, *Pantai*, and *Buaya*) from various places of Lombok Island, namely "Dusun Kerujuk, Pemenang Barat Village" and "the

forest of Pusuk Lestari, Lembah Sari village".

Maceration and Separation

500 g of each samples (*Soyun*, *Pantai*, and *Buaya*) were dried then mashed finely into powder before macerated with methanol for 3x24 hours. The crude extracts of each samples then tested using the Brine Shrimp Lethality Test (BSLT) to identify their toxicity level. The most toxic extract then chosen for the next step of separation using a vacuum liquid chromatography (VLC) technique. The stationary phase was silica gel GF60, while the mobile phase was n-heksane, DCM, methanol, etylacetate and acetone. The fractions that have the same retention factor (RF) in thin layer chromatography (KLT) were combined then were retested using BSLT to detect the most toxic fraction. Finally, the chemical compounds in the most poisonous fraction were identified applying gas chromatography-mass spectrometry (GC-MS).

Phytochemical test

Phytochemical test was conducted to analyze the compound group in the extracts qualitatively. The most toxic provenance and the combinations of the fractions from VLC was tested alkaloid, flavonoid, steroid, triterpenoid, and saponin using respective reagent.

Toxicity test using BSLT

The toxicity test was conducted using BSLT. First of all, stock solutions were made by adding 50 mg of samples into 5 mL of DMSO. The stock solutions then diluted into various concentrations, namely 10 ppm, 100 ppm, and 1000 ppm (Test Solutions). 5 mL of each test solutions were put into a tube, then added with ten (10) *Artemia salina L. Larvae*. The toxicity level was analyzed by counting larvae mortality after observed for 24 hours. The mortality percentage was counted using Abbot formula.⁹

$$\% \text{ Mortality} = \frac{(\text{sample mortality} - \text{control \% mortality})}{(100 - \text{control \% mortality})}$$

The mortality percentage then conversed into LC₅₀ using statistical analysis (Probit and regression analysis).¹⁰

$$y = ax + b$$

GC-MS Analysis

The GC-MS type is QP2010S SHIDMADZU.

In the analysis, helium was used as the carrier gas. The column type was Rtx 5 semi-polar with 30 m long, 0.25 mm diameter and 0.25 μm film thickness. Helium was used as the gas carrier with flow rate 30 mL/minute. The molecular mass range of ions was identified at 35-500m/z. the temperature program: the injection temperature was set to

260°C with initial column started at 40°C for 5 min and programmed to increase 30°C per min until it reached 260°C for 7 minutes.

RESULT AND DISCUSSION

Toxicity test

Tabel 1: Toxicity result of the methanol extract of the *G. versteegii* bark from the three provenance

Provenance	Concentration (ppm)	Log concentration	Σ larvae mortality rate after 24 h			Mortality (%)	LC ₅₀ ($\mu\text{g/mL}$)
			1	2	3		
<i>Soyun</i>	1000	3	10	10	10	100	75,86
	100	2	3	3	2	27	
	10	1	1	2	1	13	
<i>Buaya</i>	1000	3	10	10	10	100	45,94
	100	2	4	5	3	40	
	10	1	3	2	3	27	
<i>Pantai</i>	1000	3	10	10	10	100	56,36
	100	2	4	5	4	43	
	10	1	2	0	3	17	
Standard Sea Water	-	0	0	0	0	0	0
DMSO	-	0	0	0	0	0	0

Tabel 2: Toxicity result of the *G. versteegii* Buaya fractions

Fractions	Concentration (ppm)	Log concentration	Σ larvae mortality rate after 24 h			Mortality (%)	LC ₅₀ ($\mu\text{g/mL}$)
			1	2	3		
1,3,6,7,10	1000	3	10	10	10	100	489,89
	100	2	0	0	0	0	
	10	1	0	0	0	0	
2	1000	3	10	10	10	100	64,12
	100	2	9	7	7	77	
	10	1	1	0	0	3	
4	1000	3	10	10	10	100	116,81
	100	2	0	2	1	10	
	10	1	1	1	0	7	
5,9	1000	3	10	10	10	100	254,02
	100	2	1	0	0	3	
	10	1	0	0	0	0	
8	1000	3	10	10	10	100	165,20
	100	2	1	0	0	3	
	10	1	1	0	0	3	

Phytochemical test

Tabel 3: phytochemical study of the barks of *G. versteegii* Buaya

Compound class	Indication	Result
Alkaloid	White Precipitation, the solution changes into green	+
	Brown precipitation, the solution changes into brownish oranges	+
Saponin	Foam appear for \pm 10 minutes	+
Flavonoid	The solution changes into blackish green	+
Steroid	Greenish blue ring appear in the solution	+
Terpene	The solution changes into brownish red	+

GC-MS Analysis

The methanol extract has 57 compounds

identified while the fraction-2 conveys 27 compounds.




Fig. 1. GC-MS chromatogram of the *G. versteegii Buaya* extract Methanol extract b. fraction 2

DISCUSSION**Toxicity test**

The methanol extract of *G. versteegii Soyun*, *G. versteegii Buaya*, and *G. versteegii Pantai* was evaluated for the toxicity activity against Brine Shrimp Larvae. The macerate was diluted into 1000, 100 and 10 ppm and the result of the test is provided in Table 1.

Based on Table 1, the *G. versteegii Buaya* variety has shown the highest cytotoxicity with LC_{50} 45,94 $\mu\text{g}/\text{mL}$. A compound is classified as toxic if it has $LC_{50} < 1000 \mu\text{g}/\text{mL}$ according to Meyer *et al.*,⁹. Thus, *G. versteegii Buaya*, as the most poisonous variety, was chosen for the next steps that are phytochemical test and compound separation using liquid vacuum chromatography. The phytochemical test result is displayed in Table 3. Generally, the bark of *G. versteegii Buaya* is containing alkaloid, saponin, flavonoid, steroid, and terpene. Alkaloid and flavonoid have been known for their anti-cancer and

anti-tumor properties.¹¹ Hence, a further separation of the methanol extract was conducted.

The separation of the *G. versteegii Buaya* extract with liquid vacuum chromatography yielding ten fractions (F1-F10) that were obtained by adding the solution polarity with the eluen of n-hexane: ethyl acetate. The ratio were 10:0 ; 9:1 ; 7:3 ; 6:4 ; 5:5 ; 3:7 ; 1:9 respectively. These fractions then retested their toxicity level using BSLT and the result is shown in Table 2.

Based on Tabel 2, F2 has the highest toxicity activity against the *Artemia salina L. larvae* with LC_{50} 64.12 $\mu\text{g}/\text{mL}$. According to Hamidi *et al.*,¹⁰ this result is classified as forceful toxicity. If the LC_{50} is around 500-100 $\mu\text{g}/\text{mL}$, it is classified as weak, while 100-500 $\mu\text{g}/\text{mL}$ is moderate and 0-100 $\mu\text{g}/\text{mL}$ is strong. Nevertheless, the toxicity of F2 is weaker compared to the crude methanol extract of the *G. versteegii Buaya*, which have LC_{50} 45.94 $\mu\text{g}/\text{mL}$. The strong-toxicity activity is associated with the number of metabolites in the extract. Overall, these results reveal a potency of the *G. versteegii "Buaya"* as a

new anti-cancer agent. The chemical compound in the methanol extract and the F2 was identified based on GC-MS.

GC-MS Analysis

According to the literature review, overall,

five anti-cancer compounds are detected in both methanol extract and F2. Those compounds are 1,4-Benzenediol,2-methyl (1), Pyridoxylamine (2), 2,3-Dimethylhydroquinone (3), Tetramethyl-p-benzoquinone (4) and Benzofuran (5). The abundance percentage in the extracts is displayed in Table 4.

Table 4: Anticancer compounds in the methanol extract and the fraction 2 of *G. versteegii* Buaya

No	Compound	% Area in MeOH extract	% Area in Fraction 2
1	1,4-Benzenediol,2-methyl (1)	1.23	0.68
2	Pyridoxylamine (2)	1.28	5.64
3	2,3-Dimethylhydroquinone (3)	2.37	3.79
4	Tetramethyl-p-benzoquinone (4)	3.67	72.25
5	Benzofuran (5)	3.30	1.42

Tetramethyl-p-benzoquinone (4) or known as duroquinone is a part of benzoquinone compound. It is detected as the most abundance compound in both methanol extract and F2. Benzoquinone has been proved to have potential in treating breast cancer. The compound has effectively inhibiting the proliferation of estrogen receptor-positive MCF-7 cells through the NF- κ B pathway via estrogen receptor signaling.¹² Other quinone compounds also detected in the agarwood barks extracts as hydroquinone group, which are 1,4-Benzenediol,2-methyl (1) and 2,3-Dimethylhydroquinone (3). They are reported to successfully induce the death cell of A431 epidermal cell lines, SYF (Src, Yes, and Fyn), B16F10 and MDA-MB-231-cells that are implicated as key regulators of ligand-induced cellular responses including proliferation, survival, adhesion and migration. Furthermore, the process of the cell cancer inducement is rated to be safe and efficient without adverse effect.¹³

Pyridoxylamine (2), commercially known as vitamin B6, is a primary metabolite that is important for metabolism. It has a great record of its anti-cancer activity. It is reported that pyridoxylamine has suppressed cell proliferation of B16 and B16F10 murine melanoma cells and M21-HPB human melanoma cells.¹⁴ Benzofuran (5) and its derivatives like 5-Methyl-benzofurazan (6) have reported to effectively kill cancer cells line like lipoma, Hella, ovarian cancer and leukemia cells by inhibiting the cell proliferation. In addition,

benzofuran and its derivative also have antitumor activity against MCF-7 dan PC-3.¹⁵⁻¹⁹




Fig. 2. Anticancer compounds in the methanol extract and the F2 of *G. versteegii* Buaya

Other compounds that are detected in the methanol extract of *G. versteegii* Buaya are α -Caryophyllene (6) β -Santalol (7). α -Caryophyllene (6), known as humulene is sesquiterpenes that is found in the essential oil. This compound is one of the responsible compounds for the aromatic scent. Furthermore, this compound is also known for its anti-cancer activity. According to Ambroz *et al.*,²⁰, 21 Humulene is an effective treatment for intestinal, ovarian and lymphoblast cancer by inhibiting the proliferation process and oxidating effect.

β -Santalol is one of the substantial compounds in sandalwood essential oil (SEO). An exposure of SEO (2–8 μ g/mL for 24 h) is capable of breaking the single- and double-strand DNA in the MCF-7 cells, a human breast cell line. This genotoxic activity is contributed by the presence of β -Santalol, along with α -Santalol.²²




Fig. 3. Other compounds that are detected in the barks of methanol extract of *G. versteegii* Buaya

CONCLUSION

Among *G. versteegii* provenances that are found in the Lombok Island, *G. versteegii* Buaya has the most toxic effect on the brine shrimp larvae that consequently makes it as the most potential provenance to be developed as anticancer drugs.

Specifically, the methanol extract was more active in inhibiting larvae reproduction. The cytotoxicity from both methanol extract and F2 is predicted to be influenced by the same compounds, namely, 1,4-Benzenediol, 2-methyl, pyridoxylamine, 2,3-Dimethylhydroquinone, Tetramethyl-p-benzoquinone, and benzofuran. Moreover, the methanol extract has two additional compounds (α -Caryophyllene and β -Santalol) that strengthen its cytotoxic activity.

ACKNOWLEDGEMENT

We would like to give sincere gratitude to MENRISTEK (Ministry of Research and Technology) of Indonesia (contract number of research grant: 735/UN18.L1/PP/2018) for funding this research.

Conflict of Interest

The authors declare that this research has no conflict of interest.

REFERENCES

- Mulyaningsih, T.; Yamada, I. Natural resource management and socio-economic transformation under the decentralization in Indonesia: Toward Sulawesi area studies., **2007**, 365-372.
- Mulyaningsih, T.; Marsono, D.; Sumardi, S.; Yamada, I. *Jurnal Penelitian Hutan dan Konservasi Alam.*, **2017**, 14(1), 57-67.
- Mulyaningsih, T.; Marsono, D.; Sumardi; Yamada, I. *Ecology, Environment and Conservation Paper.*, **2017**, 23, 723-729.
- Setyaningrum, H. D.; Saporinto, C. *Penebar Swadaya Grup.*, **2014**.
- Hadi, S.; Muliastari, H.; Sukma, N. S.; Ratnaningsih, P. E. W. Proceeding of the 2th International Seminar on Chemistry, *Jatinangor.*, **2011**, 79-82.
- Ibrahim, A. H.; Al-Rawi, S. S.; Majid, A. M. S. A.; Rahman, N. N. A.; Abo-Salah, K. M.; Ab Kadir, M. O. *Proc. Food Scien.*, **2011**, 1, 1953-1959.
- Gunasekera, S. P.; Faircloth, G. T. *The Journal of Organic Chemistry.*, **1990**, 55(25), 6223-6225.
- Liu, J.; Wu, J.; Zhao, Y. X.; Deng, Y. Y.; Mei, W. L.; Dai, H. F. *Chinese Chemical Letters.*, **2008**, 19(8), 934-936.
- Meyer, B. N.; Ferrigni, N. R.; Putnam, J. E.; Jacobsen, L. B.; Nichols, D. E. j.; McLaughlin, J. L. *Planta medica.*, **1982**, 45(05), 31-34.
- Hamidi, M. R.; Jovanova, B.; Panovska, T. K. *Maced pharm bull.*, **2014**, 60(1), 9-18.
- Bhavana, V.; Sudharshan, S. J. S.; Madhu, D. Anticancer plants: *clinical trials and nanotechnology*, Springer., **2017**, 51-104.
- Zheng, L.; Cai, Y.; Zhou, L.; Huang, P.; Ren, X.; Zuo, A.; Meng, X.; Xu, M.; Liao, X. *International J. of Molecu. Medicn.*, **2017**, 39(1), 39-46.
- Byeon, S. E.; Yi, Y.-S.; Lee, J.; Yang, W. S.; Kim, J. H.; Kim, J.; Hong, S.; Kim, J.-H.; Cho, J. Y. *Internal. J. of Molec. Scien.*, **2018**, 19(3), 903.
- Matsuo, T.; Sadzuka, Y. *Anticancer research.*, **2019**, 39(7), 3429-3432.
- Napiórkowska, M.; Cieslak, M.; Kaomierczak-Baranska, J.; Królewska-Golinska, K.; Nawrot, B. *Molecules.*, **2019**, 24(8), 1529.
- Santha, S.; Dwivedi, C. *Anticancer research.*, **2015**, 35(6), 3137-3145.
- Shah, C. P.; Kharkar, P. S. *Journal of enzyme inhibition and medicinal chemistry.*, **2018**, 33(1), 972-977.
- Xu, J.-P. *CRC Press.*, **2016**.
- Coskun, D.; Tekin, S.; Sandal, S.; Coskun, M. F. *Journal of Chemistry.*, **2016**.
- Ambrož, M.; Boušová, I.; Skarka, A.; Hanušová, V.; Králová, V.; Matoušková, P.; Szotáková, B.; Skálová, L. *Molecules.*, **2015**, 20(8), 15343-15358.
- Ambrož, M.; Matoušková, P.; Skarka, A.; Zajdlová, M.; Žáková, K.; Skálová, L. *Molecules.*, **2017**, 22(6), 1021.
- Ortiz, C.; Morales, L.; Sastre, M.; Haskins, W. E.; Matta, J. *Evidence-Based Complementary and Alternative Medicine.*, **2016**.