



## Studies on the chemical constituents of leaves of *Phyllanthus emblica* (L.)

JAYA GUPTA, AMIT GUPTA and A.K.GUPTA

Department of Chemistry, Agra College, Agra, India.

\*Corresponding author E-mail:jayagupta6250@gmail.com

<http://dx.doi.org/10.13005/ojc/300474>

(Received: August 04, 2014; Accepted: September 26, 2014)

### ABSTRACT

A Phytochemical study was carried out on the leaves of *Phyllanthus emblica*. By using different chromatographic techniques the separation of the chemical compounds were done and structure of the compounds were elucidated by spectroscopic methods including nuclear magnetic resonance as well as mass spectrometry. Two compounds were isolated and identified; that are quercetin and  $\beta$ -sitosterol.

**Key words:** Euphorbiaceae, *Phyllanthus emblica*, quercetin,  $\beta$ -sitosterol, NMR.

### INTRODUCTION

*Phyllanthus emblica* Linn., belongs to the family Euphorbiaceae, commonly known as Indian gooseberry and amla. It is distributed in tropical and subtropical regions of India. It is excellent source of vitamin C, easily assimilated by human body<sup>1</sup>. It is found all over India, along the sea-coast districts, in deciduous forest of Madhya Pradesh. It is helpful in lowering cholesterol level<sup>2</sup> and protects from heart disease<sup>3-4</sup>, strengthens senses<sup>5</sup>, strengthens liver<sup>6</sup>. It is useful in diabetes<sup>7-8</sup>, gonorrhoea, diuretic fevers, diarrhoea<sup>9</sup>, mouth ulcers, inflammations, hair growth, headache, colic, asthma, respiratory problems. It is used as antioxidant<sup>10</sup>, aphrodisiac, antifungal, antiviral, anticancer, antigenotoxic<sup>11</sup>, antimutagenic<sup>12</sup>, chelating agent.

### EXPERIMENTAL

<sup>1</sup>H NMR and <sup>13</sup>C NMR spectra were recorded on a Bruker Advance 400 MHz spectrometer. The EI-mass was recorded on Shimadzu QP 2000 mass spectrometer. The leaves of *Phyllanthus emblica* was collected from Agra College, Agra. The leaves were air dried under shade for ten days. Then the leaves were powdered with the help of warming blender. The air dried powdered leaves (500gm) were subjected to successive hot extraction in a Soxhlet apparatus with solvents petroleum ether, ethyl alcohol and ethyl acetate. The average time period for extraction was 72 hours. The individual extracts were filtered twice and then concentrated by distillation on vacuum. The ethanolic extract (5gm) was subjected to silica gel chromatography using isopropanol-

formic acid-water (2:5:5), to give compound 1 quercetin.

A portion of the ethyl acetate extract (4 gm) was subjected to silica gel thin layer chromatography using n-hexane: acetone (80:20) solvent system to give compound 2 that is identified  $\beta$ -sitosterol. Quercetin and  $\beta$ -sitosterol were identified by comparison with data from previous NMR and mass spectra<sup>13-15</sup>.

### Quercetin

Compound (1) Slightly yellow powder; m.p 316 °C; <sup>1</sup>H NMR (400 MHz, Me OD):  $\delta$  (ppm) = 6.20 (1H, *d*, *J* = 2.0 Hz, H-6), 6.42 (1H, *d*, *J* = 2.0 Hz, H-8), 6.90 (1H, *d*, *J* = 8.2 Hz, H-5'), 7.64 (1H, *dd*, *J* = 8.3; 2.1 Hz, H-6'), 7.76 (1H, *d*, *J* = 2.1 Hz, H-2'), <sup>13</sup>C NMR (100MHz, Me OD):  $\delta$  (ppm) = 148.4 (C-2), 137.1 (C-3), 177.4 (C-4), 162.4(C-5), 99.2 (C-6), 165.6 (C-

7), 94.8 (C-8), 158.2 (C-9), 104.6 (C-10), 124.5 (C-1'), 116.2 (C-2', C-5'), 146.4 (C-3'), 150.2 (C-4'), 121.7 (C-6').

### $\beta$ -Sitosterol

Compound (2) White powder, m.p 136°C, <sup>1</sup>H NMR (CDCl<sub>3</sub>, 400MHz): 5.38(1H, dd, *J* = 5.2 Hz, H-6), 3.56(1H, tt, *J* = 11.3; 5.3 Hz, H-3), 2.34 (1H, ddd, *J* = 13.0; 5.0; 2.0 Hz, H-4a) 0.74, 0.87, 0.88, 0.89, 0.93, 1.06 (each 3H, s, MeX6). <sup>13</sup>C NMR (CDCl<sub>3</sub>, 100MHz):  $\delta$  37.1(C-1), 31.5 (C-2), 71.7(C-3), 42.2(C-4), 140.6(C-5), 121.6(C-6), 31.8 (C-7), 31.7(C-8), 50.2(C-9), 36.4(C-10), 21.2(C-11), 39.7(C-12), 42.2(C-13), 56.7(C-14), 24.2(C-15), 28.1(C-16), 56.0(C-17), 11.7(C-18), 19.3(C-19), 36.3 (C-20), 18.9(C-21), 33.8(C-22), 26.1(C-23), 45.7(C-24), 29.2(C-25), 19.7(C-26), 19.1(C-27), 23.0 (C-28), 12.1(C-29).




Fig. 1: Structure of Quercetin




Fig. 2: Structure of  $\beta$ -Sitosterol

## RESULTS AND DISCUSSION

Aim of this study was to identify and characterize the bioactive principles from the leaves of *Phyllanthus emblica*. Compound 1 is a slightly yellow powder, m.p 316 °C. The EI-mass spectrum of 1 showed the molecular ion at *m/z* 302 [M<sup>+</sup>] corresponding to the formula C<sub>15</sub>H<sub>10</sub>O<sub>7</sub> and in agreement with other spectroscopic data. The <sup>1</sup>H NMR showed that the proton of H-6 and H-8 appeared as a duplet at  $\delta$  6.20 and 6.42. The proton of H-5', H-6' and H-2' appears at  $\delta$  6.90, 7.64 and 7.76 respectively. <sup>13</sup>C NMR spectrum showed a carbonyl group at  $\delta$  177.4. The carbon bonded to hydroxyl group appeared at  $\delta$  137.1, 146.4, 150.2, 162.4 and 165.6.

Compound 2 was isolated as white powder, m.p 136°C. The EI-mass spectrum of 2 showed the molecular ion at *m/z* 414 [M<sup>+</sup>] corresponding to the molecular formula C<sub>29</sub>H<sub>50</sub>O and in agreement with other spectroscopic data. The <sup>1</sup>H NMR spectrum showed a broad triplet at  $\delta$  5.38 corresponding to H-6 olefinic proton and multiple at  $\delta$  3.56 corresponding to H-3 alpha proton six tertiary methyl singlets. <sup>13</sup>C NMR of the compound showed 29 signals for steroid skeleton which was represented by six methyl groups. The carbon bonded to the hydroxyl group C-3 appeared at 71.7. The quercetin and  $\beta$ -sitosterol has been earlier reported in various plants<sup>16-20</sup>.

**CONCLUSION**

From the physical, chemical and spectral characteristics, compound 1 and 2 were concluded as Quercetin (Fig.1) and  $\beta$ - sitosterol (Fig.2). Quercetin is flavonoid, used in asthma, eczema, hayfever, and hives. It possess anti-inflammatory, anticancer, antiviral activity, inhibit inflammatory leukotriene production<sup>21</sup>.  $\beta$ - sitosterol is phytosterol, used as antioxidant and an anti- diabetic agent. Human liver microsome studies reveals that  $\beta$ -sitosterol inhibits the cholesterol absorption, reduces the symptoms of benign prostatic

hyperplasia<sup>22</sup>, anti-inflammatory<sup>23</sup> and anti-pyretic activity. So medicinal properties of *Phyllanthus emblica* is due to the quercetin and  $\beta$ - sitosterol, furthermore scientific evaluation are required to establish therapeutic efficacy.

**ACKNOWLEDGEMENTS**

We are very grateful to University Grants Commission, New Delhi, India for their financial assistance (Grant No.F.15-39/12 (SA-II)). We are also very thankful to Dr.M.K.Rawat, Principal Agra College, Agra for their support.

**REFERENCES**

- Nisha,P.;Singhal.R.S., and Pandit, A.B.; *Int J Food Sci Nutr.* **2004**,55(5),415-422.
- Kim, H.J.; Yokozawat, Kimhy., Tohda, C.; Rao,T.P.; and Juneja,L.R.; *J Nutr Sci Vitaminol.* **2005**,51,413-418
- Yokozawa, T.; Kim, H.Y.; Kim, H.J.; Okubo, T.; Chu, D.C.; and Juneja, L.R.; *Br. J. Nutr.* **2007**,97(6)1187-1195
- Mathur,R.;Sharma,A.;Dixit,V.P.; and Verma,M.;*J Ethnopharmacol*,**1996**,50(2),61-68
- Reddy, V.D.; Padmavathi, P.V.; Kavitha, G.; Gopi, S.; and Varadacharyulu, N.,*J.Med Food* **2011**, 14(1-2),62-68
- Tasduq,S.A.;Kaisar,P.; and Gupta, D.K.; *Phytoether Res.* **2005**,19(3),193-197
- Nampoothiri,S.V.;Prathapan, A.; Cherian, O.C.; Raghu, K.G.;Venugopalan,U.V.; and Sundaresan,A.;*Food Chem Toxicol*, **2011**,49(1),125-131
- Babu,P.S.;Stainley,Mainzen,; and Prince, P.,*J. Pharm.Pharmacol*,**2004**,56,(11),1435-1442
- Mehmood,M.H.;Siddiqi,H.S.; and Gilani,A.H.; *J.Ethnopharmacol*,**2011**, 133(2),856-865
- Naik,G.H.; and Priyadarsini,K.I.;*Phytoether Res.***2005**, 19(7),582-586
- Banu,S.M.;Selevendiran,K.;Singh,J.P.; and Sakthisekaran,D.; *Hum. Exp. Toxicol*, **2004**, 23,527-531
- Madhavi, D.; Rudrama,Devi.K.; Desava, Rao. K. and Reddy, P.P.; *J. Envi. Biol.* **2007**,28,115-117
- Praveen, S.; Kader, Md.A.; Muhit, Md.A; Haque, Md.E.; and Mosaddik, Md.A.; *JAPS*,**2011** 1(9) 47-50
- Karan,S.K.;Mishra,S.K.;Pal,D.K.; and Mandal,A.;*J.Med.Plant Res.*,**2012**,6(7),1219-1223
- Trivedi, P.C.; and Chaudhary,N.; *JPR*, **2011**, 4(11),4252-4253
- Rahman,S.M.M.;Mukta,Z.A.; and Hossain,M.A.; *As J Food Ag-Ind*,**2009**, 2(1),39-43
- Akhtar,P.;Ali,M.;Sharma, M.P.;Farooqi,H.; and Khan, H.N.; *J.Phytol*, **2010**,2(3), 89-100
- Khanam,S.; and Sultana,R.; *IJSPR* **2012**,3(4), 1057-1060
- Sarin, R., and Bansal,N.; *IJRAP* **2011**, 2(3), 927-930
- Subramaniam,S.S.; and Nair, A.G.R.; *Phytochem*, **1972**,11,439
- Numata.Y.; and Tanaka,H.;*Food Chemistry*, **126**, **2011**,751-755
- Wilt,T.J.;Macdonald,R.;and Ishani,A.; *B.J.U Int.*, **1999**,83(9),976
- Loizou.S.,Lekakis,I.; Chrousos,G.P.;and Moutsatou,P.,*Mol.Nut.Food Res.*, **2010**, 54, 551.